

Extension briefs for February & March 2011

H.F. LE ROUX & J.J. BESTER

Citrus Research International

INTEGRATED PEST MANAGEMENT

Mealybug S.D. MOORE

Growers should be scouting for mealybug regularly, by inspecting underneath calyces and thereby determining percentage of fruit infested. The most effective way of doing this is to break the fruit off from the calyx. Both the fruit and the underside of the calyx should then be inspected. Where mealybug is under good biocontrol, infestation should peak during December in the northern production areas and during January in the Cape production areas. If mealybug infestation does not decline during January and February, respectively, suppression with a chemical treatment is advisable on early maturing cultivars.

Trial results have demonstrated that buprofezin (Applaud) is by far the most effective corrective option for mealybug control. It is imperative that any application of buprofezin be targeted against the younger stages of mealybug, i.e. eggs, crawlers and second instars. Where buprofezin cannot be used, methomyl and chlorpyrifos can be used but methidathion can no longer be used after petal fall. Mevinphos (at 50 ml per 100 L water) also has some effect but this is not a registered option.

This is also a good time to determine which species of mealybug are present. This is important, as it appears that the biocontrol complexes of oleander mealybug and longtailed mealybug, in particular, might not be as effective as those of citrus mealybug. Therefore, treatments can be applied more readily when either of these species is identified as the dominant species in a particular orchard. The phytosanitary status of certain species must also be borne in mind. Fortunately, recent inspections in the Eastern and Western Cape have indicated that citrus mealybug is currently the dominant species in the mealybug complex.

False codling moth S.D. MOORE

Effective false codling moth (FCM) control begins in November with diligent orchard sanitation and the application of a registered control treatment, if and where necessary. For growers in black spot-free areas who are accredited for export of fruit to the USA, it is prudent and possibly mandatory to apply a second treatment during late January or February. Growers in other areas can determine the extent of an FCM problem through weekly monitoring of traps, routine inspection of abscised fruit and the historical status of FCM levels in orchards. This will help in determining whether a follow-up treatment for FCM is necessary at this time of year. The use of the mating disruption product, Isomate, can also be considered as a stand-alone treatment or in conjunction with one or more spray treatments.

Bear in mind that the use of triflumuron (Alsystin) and teflubenzuron (Nomolt) is restricted to not later than 90% petal fall

for several important markets, making them relatively ineffective for protection of the in-season crop. Also note that the withholding period for fenpropathrin (Meothrin), which is 28 days for most markets, is 185 days for certain markets and cultivars. Cypermethrin is not allowed on fruit destined for the USA market, but is permitted in other markets with a withholding period of 28 days. Coragen (Rynaxypyr) was recently registered for FCM on citrus. However, MRLs have not yet been established for most markets, including the EU. Therefore at this stage Coragen can only be used on fruit destined for the USA and Codex A markets, with a 7-day withholding period. Delegate might be registered for FCM by the time this article appears. This will be at a higher rate than it is currently registered. However, before its use is considered, its registration status must be confirmed with the local supplier. Therefore, currently Cryptogran and Cryptex are the only pesticide sprays for FCM, which are permitted in all markets. A virus application should be applied shortly after a peak in FCM activity, determined by the use of a Lorelei pheromone dispenser in either a pipe-trap or a delta-trap. Note that due to high levels of UV irradiation in the Clanwilliam and Piketberg magisterial districts of the Western Cape and in the Northern Cape, problems may be experienced with the efficacy of virus applied in these areas during February and March. The use of Cryptogran should therefore ideally be used in these areas before the beginning of December and again after March. This problem does not exist in the rest of the country and therefore the caution is limited to the above-mentioned areas.

In addition to the insecticides, there are two mating disruption products – Isomate and Checkmate – and an attract and kill product, namely Last-Call FCM. However, all of these products are most effective when their use is initiated early in the season while FCM levels are still low. If this is not the case, initiation of their use late in the season is not recommended. Additionally, as the weather cools towards autumn, these pheromone-based products may become less effective due to a reduction in release rate. In such a case it may be necessary to follow up these treatments with a spray for FCM.

Early maturing mandarins, such as Satsumas, which will be harvested during March, should be strongly considered for a registered spray treatment for FCM during February, i.e. approximately four weeks before harvest. An effective treatment at this time should reduce post-harvest risks associated with FCM.

Bud mite T.G. GROUT

The period February to May is the optimal time for bud mite sprays and this season Mitigate (fenpyroximate) can be used during this period at 150 ml per 100 L water. The preharvest interval for most countries is 28 days but there are some minor markets that will not accept residues of this product such as Canada so it is important to check the residue restrictions. Orchards with

fruit going to markets that do not have an MRL can be sprayed immediately after removing all fruit in winter as had to be done recently with Acarol. In trials with Mitigate, this product was found to have similar efficacy to Acarol against bud mite so although a spray after harvest is not at the optimal time it will still have more impact against this pest than other unregistered options. Mitigate will also suppress citrus red mite when sprayed during autumn for bud mite and CRI research has shown that it is also effective against citrus grey mite.

Fruit fly T.G. GROUT & A. MANRAKHAN

Fruit fly monitoring and management practices should be initiated two months before the earliest expected harvest date. Recommendations remain unchanged and are as follows. Fruit flies should be controlled by use of weekly protein hydrolysate bait sprays, the use of GF120, M3 bait stations or Last Call FF. For the use of malathion in bait sprays however, there is a pre-harvest interval of 28 days. Traps must be used to indicate whether the control strategy is adequate. Growers have the option of monitoring male flies in Sensus traps with Capilure and using a threshold of 4 Medflies per week per trap, or 2 Natal flies per week per trap, or alternatively using Questlure in a Sensus trap with a threshold of 1 female fly per week per trap. If these thresholds are exceeded, additional control measures will be required.

For early maturing cultivars, particularly Satsumas, where fruit fly population pressure is usually high at the beginning of summer due to warm temperatures and presence of other highly susceptible host fruits in the vicinity, growers should use more than one mode of fruit fly control. For example, if growers are using M3 bait stations and the trap thresholds are continuously exceeded, protein bait sprays should be applied additionally until trap catches are restored below threshold levels.

Growers are strongly encouraged to set up or continue surveillance trapping for the invasive fruit fly, *Bactrocera invadens*. Surveillance trapping is conducted using bucket type traps such as Chempac Bucket trap, Moroccan trap and Lynfield trap baited with methyl eugenol. The density of traps for detection is 1 trap per km². Traps should be checked at least once a month and rebaited every 6 weeks. Trap servicing records should be kept even if no specimens are caught. Suspect specimens should be forwarded immediately to CRI for confirmation.

CROP AND FRUIT QUALITY MANAGEMENT

J.S. VERREYNNE

Internal quality: If properly timed, regulated deficit irrigation can result in an increased total soluble solids (TSS) and an increase or no response in titratable acidity. Deficit irrigation retards the breakdown of acid and can influence the solids:acid ratio at harvest for better or worse, depending on cultivar characteristics. It is mainly aimed at early cultivars like Satsuma, but other early maturing cultivars with low internal quality could benefit. No water stress should be imposed during the initial growth phase of the fruit, i.e., during and after flowering, but only during the final maturation phase, i.e., the last two months prior to harvest (January for Satsuma). Less water is applied, and at longer intervals. Therefore, irrigation is continued but at a reduced level. Trees should be irrigated lightly two weeks prior to harvest. In high rainfall areas, regulated deficit irrigation may not be successful. The deficit should

be imposed slowly, so that the trees can adjust without symptoms of drought. Severe water stress can have adverse effects on tree health and fruit quality. High nitrogen is antagonistic to the effect of deficit irrigation. Management of this technique is much easier when trees are planted on ridges and when the right scheduling equipment is used. Additionally, regulated deficit irrigation imposed the last two months prior to harvest also enhances the rate of colour development. Selective harvest of outside fruit and delaying harvest of inside fruit will result in a higher proportion of fruit with higher TSS and better colour.

Maturity indexing on early cultivars like Satsuma should commence. Maturity indexing is done to predict the rate of change in fruit maturity in order to harvest fruit at a maturity that would maintain acceptable commercial shelf life. The aim is to define changes or rate of change in acids and sugars and to build up a data base over a number of years for comparison. Random sampling of fruit every week from each of ten representative trees should start 4 to 6 weeks before the expected harvest date. Titratable acidity is determined by titration with sodium hydroxide, sugar content (Brix) is determined using a refractometer, the sugar:acid ratio calculated and fruit colour should be read from a colour chart. All the parameters mentioned above should be plotted on a graph over time. Once plotted, trends will become apparent, harvest dates can be estimated and problem areas in internal and external quality parameters can be identified and manipulated.

Fruit growth and size

Fruit growth during this time is important to achieve optimum size at harvest.

Regrowth control should be done, especially after heavy pruning earlier in the season. A lot of regrowth adversely affects fruit size and is antagonistic to fruit colour development, especially for early maturing cultivars.

Oleo incidence: Late summer vegetative growth of bearing trees should be kept to a minimum as excessive vegetative vigour during this period is associated with high incidence of oleo at harvest.

Rind colour development: Late nitrogen application and the use of heavy summer oil sprays should be avoided as these treatments are antagonistic to rind colour development.

GRONDGEDRAAGDE SIEKTES M.C. PRETORIUS

Grond en wortelmonsters behoort elke drie jaar geneem te word om sodoende die sitrusaalwurm en *phytophthora* status in sitrusboorde te bepaal. Resultate sal dien as 'n bestuurshulpmiddel wat gebruik kan word om grondpatogene effektief te beheer.

Phytophthora bruinvrot/wortelvrot

Weens die gevaar van fitotoksiteit op gevoelige sitruskultivars tydens hoë temperature, wat gedurende Februarie / Maart kan voorkom, moet die gebruik van fosfonaatblaarbespuiting **streng volgens die etiket geskied**. Hoë dag temperature, tydelike vogstremming en warm bergwinde kan veroorsaak dat fosfonate swart stippeltjies soortgelyk aan koperskade op vrugte veroorsaak. Bome moet daarom nie gespuit word as toestande nie optimaal is nie. 'n Wortelvrot beheerprogram (blaarbespuiting) sal bruinvrot ook effektief beheer. Nie een van die fosfonate is tans geregistreer teen bruinvrot nie.

Bruinvrot ontwikkel slegs wanneer die klimaatstoestand gunstig is vir die patogeen (*Phytophthora*) om te infekteer en te ontwikkel. **Indien dit dus 'n droë najaar is en geen of slegs ligte reënuitjies voorkom, is voorkomende fosfonaatblaarbespuitings nie nodig nie.** Indien dit egter 'n nat najaar is kan bome met koper of mancozeb gespuit word om bruinvrot te beheer. Hierdie twee middels is slegs voorkomend (kontakmiddels).

Bo en behalwe droogte en hitte kan 'n oormaat vogtige toestande (baie reën) ook bome onder tydelike verwelkte toestande plaas wat 'n gevaar inhou vir blaarbespuitings. Bome moet dus nie tydens of kort na sulke toestande gespuit word nie. Laastens beïnvloed drag ook 'n boom se gevoeligheid vir droogtespanning. Hoe hoër die drag, hoe gevoeliger is die boom vir uitdroging en hoe groter is die risiko vir fitotoksiteit.

Sitrusaalgewurm

Wortelmonsters kan enige tyd van die jaar getrek word om die status van die sitrusaalgewurmpopulasies in boorde te bepaal. Wyfietellings word gebruik om te bepaal of die toediening van 'n aalgewurmdoder geregverdig is. Die drempelwaarde voordat 'n aalgewurmdoder oorweeg word is 1 000 wyfies/10 g wortels. Daar word aanbeveel dat aalgewurmdodertoedienings 'n aanvang neem tydens die begin van die reënseisoen. Dit sou daarom die regte tyd wees vir produsente in die Wes-Kaap om hulle aalgewurmonsters in Maart te trek sodat hulle weet watter boorde om te behandel wanneer winterreëns begin. Residu-weerhoudingstydperke moet in ag geneem word. Dit is belangrik om 'n aalgewurmbeheerprogram te volg aangesien 'n enkele aalgewurmdodertoediening glad nie effektief is nie en het geen effek op aalgewurmpopulasies nie.

Tydens die toediening van aalgewurmdoders is dit **uiters belangrik dat ten minste 40 mm besproeiing toegedien word nadat produkte toegedien** is om te verseker dat die middels in die grondprofiel ingewas word. Met die uitsondering van Temik, loog die meeste ander aalgewurmdoders baie stadig. Die effektiwiteit van die doders word dus belemmer indien hulle nie behoorlik deur die wortelsone versprei word nie. Geen aalgewurmdoder behoort deur drupbesproeiingsstelsels toegedien te word nie. Indien toedienings in so 'n geval gedoen moet word behoort die middels as 'n bandplasing (half meter aan beide kante van die drupperlyn) oor die drupperlyn gedoen te word. Dit kan wel deur mikro-besproeiingsstelsels toegedien word.

Indien beplan word om 'n boord te verwyder behoort 'n aalgewurmmonster geneem te word voordat die boord verwyder word sodat bepaal kan word of sitrusaalgewurms teenwoordig is. Dit dien as 'n bestuursriglyn tydens 'n herplantstrategie.

CULTIVAR DEVELOPMENT

A. LEE

Growers are reminded that the Cultivar Development Division has already published a number of cultivar fact sheets on the CRI website and more will be following in the near future. This section includes an introduction to this system as well as estimated maturity charts for the different cultivars.

In addition the Rootstock section of the Production Guidelines has been updated and is now available on our website.

GEÏNTEGREERDE BEMESTING

J.G.K. COETZEE

Blaar- en grondontledings

By sitrus word blaar- en grondmonsters gedurende Februarie tot Mei geneem maar die periode kan tot Julie verleng word. Die enigste probleem met monsters wat in Julie geneem word, is die beperkte tyd om die resultate betyds te verskaf sodat met bemesting in Julie/Augustus begin kan word.

Geen ontleding hoe gesofistikeerd ookal, kan die kwaliteit van die monster verbeter nie. Bestee dus tyd, energie en aandag aan die monster. Op grond van die resultate van dié monsters word baie geld aan kunsmis bestee, terwyl die oes ook benadeel kan word. Monster elke jaar dieselfde groep bome (Indeks-bome) en hou by dieselfde prosedure en tyd. Soos by enige monster moet dit die eenheid wat gemonster word, in alle opsigte verteenwoordig.

Leaf analyses

Leaf analyses are an indicator of the nutritional status of the trees. During the research into this method a relationship was established between the concentration of the nutrient elements in the leaves and production. This relationship was developed for almost every nutrient element. For some like chloride and sodium only the maximum tolerable concentration was determined.

Let ook op die volgende spesifieke vereistes.


- Verdeel die boorde in monster eenhede, verkieslik nie groter as 5 ha nie.
- Kies twee of vier INDEKS-RYE wat die monster-eenheid in alle opsigte verteenwoordig en merk die rye.
- Gebruik elke jaar dieselfde indeksrye om blaar- en grondmonsters te neem.
- Pluk sowat 50 tot 75 blare per monster deur aan die linker- en regterkant (son- en skadukant) tussen heup- en kophoogte, blare te pluk.
- Pluk blare van agter 'n vrug, wat op dieselfde takkie as die vrug en in die lente gevorm is (Figuur 1).
- Pluk die blaarmonsters gedurende Februarie tot Mei.
- Plaas die blare in 'n skoon plastieksakkie, druk die lug uit en knoop toe.
- Merk die monsters deur 'n etiket op die sakkie te plak of aan die sakkie te bind.

Ensure that the correct leaf is picked from the trees in the index rows or blocks. The correct leaf is one that was formed during the previous spring and situated behind a fruit on the same twig as the fruit. Collect 50 to 100 leaves per sample. Take the soil sample at the same index trees. Collect 15 to 20 subsamples, mix and submit 500 g of the composite samples for analyses.

If the soil was sampled, prepared and fertilized properly before planting, it is not necessary to take leaf samples from non-bearing trees (Figuur 2). However it is never too early to monitor the nutritional status.

Grondmonsters

Grondontledings verskaf inligting wat help om te besluit watter stappe geneem kan word om tekorte, wanbalanse en oormate in die voedingstatus van die bome reg te stel. Met 'n grondontleding word gepoog om binne sekondes of minute 'n massa van die voedingstowwe uit die grond te ekstraheer, wat die bome in 8 - 10 maande sal opneem. Elke metode wat gebruik word, het dus



Figuur 1. Voorstelling van die soort blaar wat vir blaarontleding geneem moet word.




Figure 2. Illustration of the type of leaf to be sampled from non-bearing and nursery trees.




Figure 3. Taking soil samples.

net waarde indien dit gekalibreer is en presies so uitgevoer word, soos wat dit in die kalibrasie gebruik is.

Neem van grondmonsters vir bemestingsadvies

Die grondmonsters word in dieselfde indeksrye as die blaarmonsters soos volg geneem.

Monsterneming by mikrospuite

- Neem 'n submonster vanaf die oppervlak (verwyder slegs die blare, maar geen grond nie) tot 30 cm diep onder die druplyn. Gebruik 'n graaf of monsterboor.
- Neem sowat 15 tot 20 submonsters by die bome in die indeksrye. Plaas die in 'n plastiekemmer, meng deeglik en neem ± 400 g en verpak vir versending na die laboratorium.
- Merk die monster met die boordnaam (u verwysing) of boordkode plus u besonderhede.

Soil sampling at drippers (Figure 3)

- Remove the top 5 cm of soil plus debris.
- Take the sub-sample from 5 to 30 cm deep.
- Take the sample between the dripper and the perimeter of the wetted zone. If the wetted zones of two adjacent drippers overlap, take the sub-sample between the two drippers.
- Collect 15 to 20 sub-samples at the index trees. Put the sub-samples in a plastic bucket, mix properly and retain ± 500 g for sending to the laboratory.
- Mark the samples with your name and that of the orchard plus all relevant information on a label and stick or tie it to the outside of the container.

POST HARVEST PATHOLOGY

Waste prevention checklist

K.H. LESAR

Sanitation: NB: For reducing fungal spore load, as well as keeping FCM and fruit fly under control – remove all fallen fruit and decayed fruit from the orchard. Bury or macerate fallen fruit and allow to dry in the sun **away from the orchards**.

Remove dead wood from all citrus trees to reduce the spore load of the latent citrus pathogens.

Good fruit fly control: Use traps and bait regularly.

FCM: Apply pre-harvest sprays where trap counts and fruit drop due to FCM is high.

Skirt trees for brown rot control: Ensure that trees are adequately skirted, preventing low hanging fruit, especially in heavily laden trees, and thereby reducing the risk of *Phytophthora* brown rot infection during the rainy season.

Do not pack any fallen fruit that could be infected.

Swaar reënval - *Phytophthora* bruinvrot waarskuwing!!!!

Phytophthora bruinvrot word versprei wanneer *Phytophthora nicotianae* of *P. citrophthora* spore tydens reën vanaf besmette grond op vrugte spat. Infeksie vind plaas en die vrug vrot na 'n tydperk van 4-6 dae. Dit gebeur dus dat geïnfecteerde vrugte gepluk en gepak kan word en tydens versending bruinvrot kan ontwikkel. 'n Enkele vrot vrug kan die hele karton vrugte besmet.

Die swam penetreer die skil binne drie ure en is dan buite bereik van die swamdoders wat in die pakhuis gebruik word soos TBZ en Imazalil. Dit is dus

belangrik dat bruinvrot voorkomend in die boord, voor pluk, behandel word.

Prevent injuries:

Test for injuries. "Indigo-Carmine" can be used for this purpose. Test both in the orchard and the packhouse. **NB: Ensure proper picking practices are adopted. There are far too many injuries every season, resulting in abnormal waste levels.**

Let Wel: Sorg dat gepaste plukpraktyke toegepas word. Te veel plukbeserings het elke seisoen onreëlmatige vlakke van bederf veroorsaak.

Apply packhouse fungicides with care: Check the mixing/application rates.

Packhouse sanitation:

- Never allow any fruit to lie around in the packhouse and develop spores.
- Constantly monitor concentrations of sanitisers in dump tanks, de-scaler water, rinses, etc.
- Spray the packhouse out with sanitisers regularly and immediately after finding a single mouldy fruit.
- Spray out trailers/picking bins with a suitable sanitiser before they leave for the orchard.
- Transport cartons to the ports as soon as possible and prevent packed fruit standing on the packhouse floor where it is hot. Green mould develops faster at 10°C than at 4.5°C.
- Store retention samples for each consignment and check regularly for waste and other developing factors.

The control of post-harvest diseases on export citrus using the post-harvest fungicide thiabendazole

There seems to be a general reduction in the use of thiabendazole within the SA citrus industry. This is particularly alarming since latent pathogen infections have been observed in abundance during recent citrus production seasons.

Why use thiabendazole (TBZ)? TBZ was the first fungicide registered (1960s) for the control of the *Penicillium* moulds and the latent pathogens, *Diplodia* stem-end rot, *Phomopsis* stem-end rot and *Anthraco*se on citrus fruit. TBZ and benomyl belong to the benzimidazole group of fungicides.

The benzimidazoles are distinguished from other traditional fungicides in that they control diseases both by contact and systemic action. Due to the extensive pre-harvest application of benomyl for the control of citrus black spot and the post-harvest application of TBZ for control of the *Penicillium* moulds, populations of *Penicillium* moulds that were resistant to the benzimidazoles developed rapidly. There is therefore an unfortunate perception in the industry that TBZ is of no value in controlling important post-harvest pathogens.

However, TBZ is still effective in controlling the latent pathogens on citrus: *Diplodia* stem-end rot, *Phomopsis* stem-end rot and *Anthraco*se. All export citrus should therefore be treated with TBZ.

Application of TBZ in the wax to fruit also reduces the risk of some rind conditions developing during storage and export, e.g. chilling injury, pitting etc.

Some guidelines for reducing the risk of chilling injury on

grapefruit exported under extended cold sterilisation conditions: Citrus in general is known to be sensitive to cold damage (chilling injury) during shipping and storage, but certain cultivars (some soft citrus cultivars, lemons and grapefruit varieties), are particularly prone to chilling injury, especially when exposed to "cold sterilisation" temperatures. It is especially the yellow pigmented citrus cultivars viz. lemons, Marsh grapefruit, and even the yellow areas of Star Ruby and Rose grapefruit which are the most sensitive, as they do not contain the carotenoids which act as anti-oxidants that protect the fruit against chilling injury.

The extended cold sterilisation treatment, as recently adopted by China (24d at -0.6°C), is particularly problematic. It is generally accepted that it is not feasible to export lemons under these conditions. However, whereas grapefruit is also highly sensitive to chilling injury, there are some practices, as summarised in this document, that can be followed to reduce the risk of chilling injury.

Picking window

The South African grapefruit season, in the traditional production areas, extends from the middle of March to the end of June. The picking window for grapefruit is often manipulated in an attempt to access markets early or to extend the season. However, harvesting grapefruit too early in

CONTINUED ON PAGE 32



**MED
AUTOMATION**

28 Donkersvliet Street • PO Box 3014 Daljosaphat Paarl 7620
Tel: 021 868 1652 • Fax: 021 868 1739

Quality machine builders

**The complete solution for all
packing and sorting of fruit and vegetables**







the season, when the fruit rind is still “immature” and also at or beyond the end of the season when the fruit is well coloured and “very mature”, is when grapefruit is most sensitive to cold injury. It is a major risk to export such sensitive fruit to markets where cold sterilisation is a requirement.

Thorough maturity indexing is essential to determine the ideal harvesting window. Commencing 5 weeks before anticipated harvest, pick samples of grapefruit (20-25 fruit). Mark the representative trees (data trees from different rootstocks, selections, tree ages or microclimates). Evaluate and record average fruit colour and full internal quality assessments. Repeat every week until optimal harvest date, ensuring that the samples are drawn similarly for comparison. Plot the results on a graph to determine whether the season is early or late compared to the previous year, thereby determining the optimal picking window for the specific cultivar.

Commencement of export packing of grapefruit to “cold steri” markets should start 14 days later, as the rinds will still be too cold sensitive at the beginning of the normal optimal picking window. Harvesting of grapefruit for such “cold-steri” markets should also not be extended beyond the end of the optimal picking window.

Post-harvest wilt conditioning

Conditioning (wilting) trials where Marsh grapefruit (exported to Japan) was “conditioned” for 2, 4 and 6 days at 16°C and 20°C prior to cold treatment, showed a dramatic reduction in the incidence of chilling injury relative to the non-conditioned control fruit. Unfortunately, extending the time between packing and introduction into the cold chain can also increase the incidence of post-harvest rind pitting and decay in sensitive fruit. Nonetheless, wilting at ambient for 7 days is part of the standard handling procedure for grapefruit exported to Japan and should be implemented by anybody wishing to risk exporting grapefruit under an extended cold treatment regime.

The role of TBZ

It is known that inclusion of thiabendazole (TBZ) in citrus wax applied to grapefruit can significantly reduce the incidence of chilling injury. Inclusion of TBZ is also a good standard packhouse procedure and should be used if anybody wishes to risk exporting grapefruit under these extended cold sterilisation conditions.

Pre-cooling and Storage

A critical factor affecting the extent of chilling injury of grapefruit, is the duration of exposure to temperatures below 4.5°C. This exposure period is cumulative and can occur during pre-cooling (the period prior to loading during which the temperature of the fruit is reduced to the cold sterilisation level), the cold sterilisation treatment itself and post-shipping storage. Pre-cooling for 3d is a compulsory component of the disinfestation treatment, but any other pre-loading storage at temperatures below 4.5°C should be avoided. Storage of grapefruit after shipping should be at an “intermediate storage temperature” of 7 to 8°C and should be kept to the minimum necessary duration.

The role of waxes

Why wax fruit? The main purpose of a wax emulsion is to:

- Protect fruit against moisture loss, which results in longer shelf

life and less weight loss of the fruit.

- To provide **shine** to the fruit at the point of sale. It is important to note that this shine needs to be sustained throughout the chain of distribution (some waxes break down more rapidly than others).

- Provide a **barrier of protection against chilling injury and fungi**.

The type of wax that you choose and the way in which you apply the wax will have a significant influence on the above points.

Choosing the right wax: The “lighter” waxes (e.g. lower solids or carnauba based waxes) offer less resistance to gas transfer (respiration) than “heavier” waxes (e.g. shellac or high solids polyethylene based waxes). Thus light wax emulsions will protect fruit with sensitive rinds far better than a heavier wax emulsion and will also allow for better colour development. On the other hand lighter waxes break down more quickly, and are therefore not suitable for long storage programmes, especially cold sterilization programmes. Research has indicated that “heavy waxes” that slow breathing (respiration) down and retain a high level of CO₂ (10%+) on the surface of the fruit, reduce the incidence of CI. Unfortunately the use of heavy waxes may increase the incidence of post-harvest rind pitting in sensitive fruit. Nonetheless, the high risk of chilling injury on grapefruit under conditions of extended sterilisation, make it appropriate to consider preferentially using such waxes when exporting to markets that require such extended cold treatment.

Wax application

Uneven waxing, under-waxing and over-waxing all have a deleterious effect on fruit quality. The wax barrier need not be thick, and in fact a very thin barrier provides enough of an integral film to prevent most of the moisture loss without interfering with the respiration process. NB: Adhere to the wax manufacturer’s recommended application rate – **do not apply your own rate!!! Adhere to the product label instructions.**

Please note: • Fruit should be dry before waxing. Where possible the packhouse should use a hot water bath for fungicide application as this helps the drying of fruit before waxing.

- Fruit must move evenly through the waxing unit and flow of fruit entering the packing line must be consistent.
- When fruit leaves the waxing unit, all parts of the fruit needs to be covered with a film of wax. Examine fruit after the waxing unit regularly.
- Brushes in the wax applicator must be in good condition and should rotate at a speed of about 90 rpm. The last brush in the wax applicator should always be wet, so as not to remove wax from the fruit.

Over application

- Inhibits the breathing (respiration) of the fruit.
- The movement of oxygen and CO₂ on the surface of the fruit is inhibited resulting in poor colour development and off-flavours caused by the process of an aerobic fermentation.
- Inhibits colour break.
- Encourages rind disorders on sensitive fruit.
- Unnecessary expense!!

Under application

- Excessive weight loss and shrinkage of fruit.
- Poor shelf life / storage.
- Susceptibility to chilling injury.